

A new method to improve photometric redshift reconstruction

Applications to the Large Synoptic Survey Telescope

Alexia Gorecki¹, Alexandra Abate^{2,3}, Réza Ansari², Aurélien Barrau¹, Sylvain Baumont¹, Marc Moniez² and Jean-Stephane Ricol¹

¹ Laboratoire de Physique Subatomique et de Cosmologie, UJF/INP/CNRS/IN2P3, 53 avenue des Martyrs, 38026 Grenoble cedex, France

² Laboratoire de l'Accélérateur Linéaire, Univ. Paris Sud/CNRS/IN2P3, Bat. 200, 91898 Orsay cedex, France

³ Physics Department, University of Arizona, 1118 East 4th Street, Tucson, AZ 85721, USA
e-mail: abate@email.arizona.edu

Received ???; accepted ???

ABSTRACT

Context. In the next decade the Large Synoptic Survey Telescope will become a major facility for the astronomical community. However accurately determining the redshifts of the observed galaxies without using spectroscopy is a major challenge.

Aims. Reconstruction of the redshifts with high resolution and well-understood uncertainties is mandatory for many science goals, including the study of baryonic acoustic oscillations (BAO). We investigate different approaches to establish the accuracy that can be reached by the LSST six-band photometry.

Methods. We construct a realistic mock galaxy catalog, based on the GOODS survey luminosity function, by simulating the expected apparent magnitude distribution for the LSST. To reconstruct the photometric redshifts (photo-z's) we consider a template-fitting method and a neural network method. The photo-z reconstruction from both of these methods is tested on real CFHTLS data and also on simulated catalogs. We describe a new technique to efficiently remove catastrophic outliers via a likelihood ratio statistical test that uses the posterior probability functions of the fit parameters and the colors.

Results. We show that the photometric redshift accuracy will meet the stringent LSST requirements up to redshift ~ 2.5 after a selection based on the likelihood ratio test or a selection based on the apparent magnitude, for galaxies with $S/N > 5$ in at least 5 bands. The former selection has the advantage of retaining roughly 35% more galaxies for a similar photo-z performance compared to the latter. Photo-z reconstruction using a neural network algorithm is also described. In addition we utilize the CFHTLS spectro-photometric catalog to outline the possibility of combining the neural network and template-fitting methods.

Conclusions. We demonstrate that the photometric redshifts will be accurately estimated with the LSST if a Bayesian prior probability and a calibration sample are used.

Key words. cosmology – photometric redshift – large scale survey – LSST – CFHTLS

1. Introduction

The Large Synoptic Survey Telescope (LSST) has an optimal design for investigating the mysterious dark energy. With its large field of view, its short exposure time and its high transmission bandpasses, the LSST will be able to observe a tremendous amount of galaxies, out to high redshift, over the visible sky from Cerro Pachón over ten years. This will lead to an unprecedented study of dark energy, among other science programs such as the study of the Milky Way and our Solar System (LSST Science Collaboration 2009).

One of the main systematic uncertainties in the cosmological analysis will be tightly related to errors in the photometric redshift (photo-z) estimation. Estimating the redshift from the photometry alone (Baum 1962) is indeed much less reliable than using spectroscopy, although it does allow measurements to be obtained for vastly more galaxies, in particular those that are very faint and distant. Photo-z estimates are mainly sensitive to characteristic changes in the galaxy's spectral energy distribution (SED) such as the Lyman and the Balmer breaks at 100nm and 400nm, respectively. Incorrect identifications between these two main features greatly impact the photometric redshifts, giving rise to catastrophic outliers. Mis-characterizing the propor-

tion of such outliers will strongly impact the level of systematic uncertainties.

There are basically two different techniques to compute the photo-z. On the one hand, *template-fitting methods* (e.g. Puschell et al. 1982; Bolzonella et al. 2000) fit a model galaxy SED to the photometric data and aim to identify the spectral type, the redshift and possibly other characteristics of the galaxy. It has been proven that using spectroscopic information in the template-fitting procedure, for example by introducing a Bayesian prior probability (Benitez 2000) or by modifying the SED template (Budavári et al. 2000; Ilbert et al. 2006) improves the photo-z quality. This highlights the necessity to have access to at least some spectroscopic data.

On the other hand, *empirical methods* extract information from a spectroscopic training sample and are therefore generally limited to the spectroscopic redshift range of the sample itself. Among these, the empirical color-redshift relation (Connolly et al. 1995; Sheldon et al. 2012) and neural networks (Vanzella et al. 2004; Collister & Lahav 2004) are commonly used.

In this paper, we address the issue of estimating the photo-z quality with a survey like the LSST, and in particular, we introduce a new method that aims to remove most of the galaxies

with catastrophic redshift determination (hereafter called outliers). We utilize a galaxy photometric catalog simulated for a study of the uncertainty expected from the LSST on the dark energy equation of state parameter from baryon acoustic oscillations (BAO). The related results will be presented in a companion paper by Abate *et al.* *in prep.*

Based on a Bayesian χ^2 template-fitting method, our photo- z reconstruction algorithm gives access to the posterior probability density functions (pdf) of the fit parameters. Using a training sample, the *likelihood ratio* test, based on the characteristics of the posterior pdf and the colors, is calibrated and then applied to each galaxy in the photometric sample. The technique is tested on a spectro-photometric catalog from the T0005 data release of CFHTLS¹ matched with spectroscopic catalogs from VVDS (Le Fevre *et al.* 2004; Garilli *et al.* 2008), DEEP2 (Newman *et al.* 2012) and zCOSMOS (Lilly *et al.* 2007). We also outline the possibility to discard outlier galaxies by using photometric redshifts estimated both from the template-fitting method and from a neural network.

Finally we illustrate the modification to the systematic and statistical uncertainties on the photo- z when the redshift distribution of the training sample is biased compared to the actual redshift distribution of the photometric catalog to be analyzed.

The paper is organized as follows. The LSST is presented in section 2, followed by our simulation method in section 3. In the latter, the simulation steps employed and the physical ingredients required to produce the mock galaxy catalogs are described. In section 4, the mock galaxy catalogs for GOODS, CFHTLS and LSST are presented and validated against data for the former two surveys. Our template-fitting method and the likelihood ratio test are described in section 5. The performance of our photo- z template-fitting method is shown in section 6. In section 7, the photo- z neural network technique and its performance in conjunction with our template-fitting method is investigated. Finally, we give a brief discussion of the current limitations of our simulations in section 8, and conclude in section 9.

Throughout the paper, we assume a flat cosmological Λ CDM model with the following parameter values: $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$, $\Omega_k = 0$, $H_0 = 70$ km/s/Mpc.

2. The Large Synoptic Survey Telescope

The LSST is a ground-based optical telescope survey designed, in part, to study the nature of dark energy. It should be one of the fastest and widest telescopes of the coming decades. The same data sample will be used to study the four major probes of dark energy cosmology: type 1a supernovae, weak gravitational lensing, galaxy cluster counts and baryon acoustic oscillations.

It will provide unprecedented photometric accuracy with six filters (u, g, r, i, z, y). In the nominal design each filter is expected to have a transmission higher than 95% in the bandpass and lower than 0.01% outside, with very sharp edges. Moreover, the LSST CCDs are expected to have a high quantum efficiency compared to current detectors. The LSST will be a modified

Table 1. Number of visits and 5σ limiting apparent magnitudes, for one year and ten years of LSST operation (Ivezic *et al.* 2008; Petry *et al.* 2012).

One year of observation						
	u	g	r	i	z	y
Number of visits	5	8	18	18	16	16
Limiting apparent magnitude	24.9	26.2	26.4	25.7	25.0	23.7
Ten years of observation						
	u	g	r	i	z	y
Number of visits	56	80	184	184	160	160
Limiting apparent magnitude	26.1	27.4	27.5	26.8	26.1	24.9

Table 2. LSST photo- z requirements for the high signal-to-noise “gold sample” subset with $i < 25.3$. The parameters are defined as follows: $\sigma_z/(1+z)$ is the root-mean-square scatter in photo- z ; η is the fraction of 3σ outliers at all redshifts; e_z is the bias, defined as the mean of $(z_p - z)/(1+z)$ at a given z , where z_p is the photo- z .

quantity	requirement	goal
$\sigma_z/(1+z)$	< 0.05	< 0.02
η	$< 10\%$	
$ e_z $	< 0.003	

Table 3. Number of galaxies that are both observed photometrically with the CFHTLS survey and spectroscopically.

Spectroscopic survey	CFHTLS field	No. of galaxies	Refs.
VVDS Deep1	W1 & D1	2011	(1)
DEEP2 Data Release 3	W3 & D3	5483	(2)
VVDS f22	W4	4485	(3)
zCOSMOS	D2	2289	(4)

References. (1) Le Fevre *et al.* (2004); (2) (Newman *et al.* 2012); (3) Garilli *et al.* (2008); (4) Lilly *et al.* (2007).

Paul-Baker telescope with three mirrors among which the primary mirror measures 8.4 m diameter. The camera at the center of the tertiary mirror is composed of a field corrector made of three lenses, a carousel of five filters with an additional sixth filter and a focal plane array of 63.4cm diameter. The latter is composed of 189 science CCDs of 4096×4096 10 μ m pixels, cooled down by a cryostat. The field of view of the LSST will be 9.6 deg² and the survey should cover 30 000 deg² of sky visible from Cerro Pachón.

The LSST will perform two back-to-back exposures of 2×15 sec with a readout time of 2×2 sec. The number of visits and the 5σ limiting apparent magnitude in each band, for one year and ten years of the running survey are listed in table 1. With such deep observations, photometric redshifts will necessarily be computed in an essentially unexplored redshift range.

The photo- z requirements as published in the LSST Science Book (LSST Science Collaboration 2009) are given in table 2.

The final specifications of the LSST are subject to change; see Ivezic *et al.* (2008) for the latest numbers.

¹ Based on observations obtained with MegaPrime/MegaCam, a joint project of CFHT and CEA/DAPNIA, at the Canada-France-Hawaii Telescope (CFHT), which is operated by the National Research Council (NRC) of Canada, the Institut National des Science de l’Univers of the Centre National de la Recherche Scientifique (CNRS) of France, and the University of Hawaii. This work is based in part on data products produced at TERAPIX and the Canadian Astronomy Data Centre as part of the Canada-France-Hawaii Telescope Legacy Survey, a collaborative project of NRC and CNRS.




Fig. 1. SED templates linearly interpolated from the original six templates from Coleman et al. (1980) and Kinney et al. (1996). The original templates are drawn in red.

3. Simulation of galaxy catalogs

This section begins by describing the main steps carried out in order to produce a catalog of galaxy photometric data. The rest of the section describes the ingredients used to generate these catalogs in more detail.

3.1. Simulation steps

3.1.1. Simulating galaxy distributions

In order to simulate the galaxy catalog, we first compute the total number of galaxies N within our survey volume between absolute magnitudes M_1 and M_2 . Then for each of these N galaxies, we assign redshifts and galaxy types.

If ϕ is the sum of luminosity functions over the *Early*, *Late* and *Starburst* galaxy types (see section 3.2.1 for more details), then the number of galaxies N_g is given by

$$N_g = \int_0^6 \int_{M_1}^{M_2} \phi(M, z) H_0 (1+z)^2 d_A(z)^2 E(z)^{-1} \Omega dz dM,$$

where M is the absolute magnitude in some band, $d_A(z)$ is the angular diameter distance, the function $E(z)$ is defined by

$$E(z) = \sqrt{\Omega_m (1+z)^3 + \Omega_\Lambda (1+z)},$$

and Ω (no subscript) is the solid angle of the simulated survey. The redshift range is chosen so as not to miss objects that may be observable by the survey. The exact choice of M_1 and M_2 is not critical because (i) at the bright limit the luminosity function goes quickly to zero; therefore as long as M_1 is less than -24 the integral is not affected, and (ii) as long as M_2 is chosen to be fainter than the maximum absolute magnitude observable by the survey, then all galaxies that are possible to observe will be included; we chose $M_2 = -13$. The redshift z_s of each simulated galaxy is drawn from the cumulative density function:

$$C_z(z_s) = \frac{\int_0^{z_s} \int_{M_1}^{M_2} \phi(M, z') H_0 (1+z')^2 d_A(z')^2 E(z')^{-1} dz' dM}{\int_0^6 \int_{M_1}^{M_2} \phi(M, z) H_0 (1+z)^2 d_A(z)^2 E(z)^{-1} dz dM}.$$




Fig. 2. Extinction by dust as a function of λ for three different color excess values: $E(B-V) = 0.1$ in black, $E(B-V) = 0.2$ in red and $E(B-V) = 0.3$ in green. The solid lines correspond to the Cardelli law (Cardelli et al. 1989). The bump around 217 nm can be explained by graphite or the presence of Polycyclic Aromatic Hydrocarbons (cf for example Clayton et al. 2003). The dashed lines correspond to the Calzetti law (Calzetti et al. 2000).

Once the redshift of the galaxy, denoted by z_s , is assigned, the absolute magnitude M is drawn from the following cumulative density function

$$C_M(M, z_s) = \frac{\int_{M_1}^M \phi(M', z_s) dM'}{\int_{M_1}^{M_2} \phi(M', z_s) dM'}.$$

Finally, a broad galaxy type is assigned from the observed distribution of each type at redshift z_s and absolute magnitude M . This distribution is constructed from the type-dependent luminosity functions. Therefore each galaxy is designated a broad type value of either: *Early*, *Late* or *Starburst*. An SED from the SED library is then selected for each galaxy, according to the simulation procedure described in section 3.2.2.

3.1.2. Simulating the photometric data

The simulated apparent magnitude $m_{X,s}^{[12]}$ in any LSST band X , with transmission $X(\lambda)$, for a galaxy of SED type $T_s^{[13]}$, redshift z_s , color excess $E(B-V)_s$ and absolute magnitude M_s , is generated as follows:

$$m_{X,s} = M_s + \mu(z_s) + K_{XY}(z_s, T_s, E(B-V)_s),$$

where $\mu(z_s)$ is the distance modulus and $K_{XY}(z_s, T_s, E(B-V)_s)$ is the K-correction defined as described in Hogg (1999) for spectral type T_s , with flux observed in observation-frame band X and M_s in rest-frame band Y . It is expressed as

$$K_{XY}(z_s, T_s, E(B-V)_s) = -2.5 \log \left[\frac{1}{1+z_s} \frac{\int d\lambda \lambda T_s \left(\frac{\lambda}{1+z_s}, E(B-V)_s \right) X(\lambda) \int \frac{d\lambda}{\lambda} Y(\lambda)}{\int \frac{d\lambda}{\lambda} X(\lambda) \int d\lambda \lambda T_s(\lambda, E(B-V)_s) Y(\lambda)} \right].$$

Then, the corresponding simulated flux $F_{X,s}$ is computed as

$$F_{X,s} = F_{X,0} 10^{-0.4m_{X,s}},$$

where

$$F_{X,0} = c \int X(\lambda) / \lambda d\lambda.$$

² Subscript s stands for the simulated value.

³ Here type T_s refers to the actual SED of the galaxy, and not the broad type value e.g. *Early*, *Late* or *Starburst*.




Fig. 3. LSST transmission curves shown by the solid lines and CFHTLS transmissions shown by the dashed lines. The transmission takes into account the transmission of the filter itself, the expected CCD quantum efficiency, the telescope optical throughput and the mean atmospheric transmission.

c is a constant that depends on the luminosity of the galaxy, the units the flux is measured in and the zeropoint of the AB magnitude system. The uncertainty on the flux is then given by

$$\sigma(F_{X,s}) = 0.4\sigma_X(m_{X,s})F_{X,s}\ln(10).$$

The simulated observed flux $F_{X,obs}$ is drawn from a Gaussian with mean $F_{X,s}$ and standard deviation $\sigma(F_{X,s})$. This is correct as long as the flux is large enough to be well distributed with a Gaussian distribution. The uncertainty $\sigma_X(m_{X,s})$ on true magnitude in band X is given by Eq. 2 in section 3.2.5.

Note that the apparent magnitude uncertainty $\sigma_X(m_{X,s})$ depends on the number of visits $N_{X,vis}$. We have performed the simulation for two sets of values of $N_{X,vis}$ that correspond to one and ten years of observations with the LSST, according to the $N_{X,vis}$ given in table 1.

Throughout the paper the quantity z_s refers to the simulated or “true” value of the redshift. Here we also assume that a spectroscopic redshift obtained for one of the simulated galaxies has a value equal to z_s . Therefore the value z_s can be also considered to be the galaxy’s spectroscopic redshift with negligible error.

3.2. Ingredients

3.2.1. Luminosity function

The luminosity function describes probabilistically the expected number of galaxies per unit volume and per absolute magnitude. If the luminosity functions are redshift- and type-dependent, then they give the relative amount of galaxies for each galaxy type at a given redshift.

We use luminosity functions measured from the GOODS survey (Dahlen et al. 2005). The luminosity functions here are modeled by a parametric Schechter function that takes the form:

$$\phi(M) = 0.4\ln(10)\phi^*y^{(\alpha+1)}\exp(-y)$$

$$y = 10^{-0.4(M-M_*)}$$

where M is the absolute magnitude in the B-band of GOODS WFI, and M_* , ϕ^* , and α are the parameters defining the function. Their values can be obtained from Dahlen et al. (2005).




Fig. 4. Histograms of the apparent magnitude in the R band comparing the GOODS simulated data (black points with errorbars) to the actual GOODS data (red stars).

3.2.2. Spectral Energy Density (SED) library

We built an SED library composed of 51 SEDs. They were created by interpolating between six template SEDs, described here:

- the *Early*-type El, the *Late*-types Sbc, Scd and the *Starburst*-type Im from Coleman et al. (1980),
- the *Starburst*-types SB3 and SB2 from Kinney et al. (1996).

These six original SEDs were linearly extrapolated into the UV by using the GISEL synthetic models from Bruzual & Charlot (2003). The interpolated spectra of the 51 types are displayed in Fig 1. In the following, we denote by T_s the true spectral type (SED) of the galaxy.

Each galaxy is assigned an SED from this library using a flat probability distribution based on their broad type value, originally assigned as either *Early*, *Late* or *Starburst* (see section 3.1.1). This way of generating the spectra may not be as optimal as using more realistic synthetic spectra, but it has heuristic advantages. For example, there is an easy way to relate the luminosity function main galaxy type to an SED type, and additionally it is much faster, in terms of computing time, to produce a large amount of galaxy spectra at different evolutionary stages. We are aware, however, that this linear interpolation may bias photometric redshifts estimated using a template-fitting method, because real galaxies are probably not evenly distributed across spectrum space. Therefore this feature may allow the neural network method to be more effective in estimating the redshift.

3.2.3. Attenuation by dust and Intergalactic Medium (IGM)

The reddening caused by dust within the target galaxy is quantified in our simulation by the color excess term $E(B-V)$. Together with this term, the Cardelli law (Cardelli et al. 1989) is used for the galaxies closest to the El, Sbc, and Scd spectral types, whereas the Calzetti law is used for the galaxies closest to the Im, SB3, and SB2 spectral types. In Fig. 2, the reddening laws are plotted for three values of the color excess. The parameter $E(B-V)$ is drawn from a uniform distribution between 0 to 0.3 for all galaxies, except for galaxies closest to the El type. Elliptical galaxies are composed of old stars and contain little or no dust, therefore $E(B-V)$ is drawn only between 0 to 0.1 for these galaxies.

Another process to be considered is the absorption due to the intergalactic medium (IGM). It is caused by clumps of neu-




Fig. 5. Histograms of the $b - r$ term for different apparent magnitude ranges. Left-hand panel ($20 < r < 22$) corresponds to the bright galaxies and right-hand panel ($24 < r < 25.5$) corresponds to the very faint galaxies. The solid black lines correspond to the simulation and the solid red lines correspond to the GOODS data. The dotted colored lines correspond to the main spectral types in the simulation.

tral hydrogen along the line of sight, and is well-modeled by the Madau law (Madau 1995). As the absorption occurs at a fixed wavelength in the hydrogen reference frame, it is redshift-dependent in the observer frame. Strong features in the optical part of the SEDs are induced by the IGM at redshifts above about $z \sim 2.8$ in the LSST filter set, when the Lyman- α forest has shifted into the LSST band passes. Here we assume this absorption to be constant with the line of sight to the galaxy. An investigation of the effect of the stochasticity of the IGM will be the subject of future work.

3.2.4. Filters

The six LSST bandpasses displayed in Fig. 3 take into account the quantum efficiency of the CCD, the mean atmospheric transmission, the filter transmission, and the telescope optical throughput. The CFHTLS filter set⁴ is also displayed in the same figure. We expect to be able to obtain good photo- z estimates up to redshifts of about 1.2 for CFHTLS and 1.4 for LSST when the 4000 Å break is redshifted out of the filter sets. At redshifts above 2.5 the precision should improve dramatically when the Lyman break begins to redshift into the u -band.

3.2.5. Apparent magnitude uncertainties for the LSST

The apparent magnitude uncertainties for the LSST are computed following the semi-analytical expression from the LSST Science Book (LSST Science Collaboration 2009). This expression has been evaluated from the LSST exposure time calculator, that takes into account the instrumental throughput, the atmospheric transmission, the airmass among other physical parameters.

The random error on the (true) magnitude m_X (subscript s has temporarily been dropped) can be written as

$$\sigma_{rand,X}^2 = (0.04 - \gamma_X)x + \gamma_X x^2, \quad (1)$$

where the value of γ_X is reported in the LSST Science Book for each filter X . The variable x is defined as follows:

$$x = 10^{0.4(m_X - m_{5,X})},$$

⁴ The CFHTLS transmissions have been downloaded from <http://www1.cadc-ccda.hia-ihp.nrc-cnrc.gc.ca/community/CFHTLS-SG/docs/extra/filters.html>.




Fig. 6. Redshift distribution of the spectroscopic sample for the different CFHTLS fields. The histograms are stacked.

where $m_{5,X}$ is the 5σ limiting apparent magnitude in the X band, such that $\sigma(m_{5,X}) = 0.2$, and is given by the following expression

$$m_{5,X} = C_X + 0.5(m_X^{sky} - 21) + 2.5 \log(0.7/\theta_X^{see}) + 1.25 \log(t_v/30) - k_X(A - 1),$$

where the parameter C_X accounts for the total instrumental throughput; m_X^{sky} is the sky background apparent magnitude; θ_X^{see} is the seeing; t_v is the exposure time in seconds; k_X is the atmospheric extinction; A is the air mass. All of these parameters are reported in the LSST Science Book for median observing conditions.

The total uncertainty on the apparent magnitude includes a systematic uncertainty that comes from the calibration, such that the photometric error in band X is

$$\sigma_X(m_X) = \sqrt{\sigma_{rand,X}^2 + \sigma_{sys,X}^2}, \quad (2)$$

where $\sigma_{sys,X}$ is taken to be equal to 0.005 and is the photometric systematic uncertainty of the LSST for a point source. We have adopted this simple formula defined for point sources, and have used it for extended sources. A more realistic computation of this uncertainty for extended sources will be completed in future work.




Fig. 7. Evolution of the colors $u - g$ and $i - z$ as a function of redshift computed from the K-correction (colored lines) and measured from the CFHTLS data (black dots). The departure of the computed colors from the data is not significant since not all the galaxy types exist in significant numbers within the CFHTLS data at all redshifts.




Fig. 8. *Left-hand side:* Cumulative distribution of the apparent magnitude in the i band for the LSST simulation compared to the SDSS measurement from the “Stripe 82” region (cf Abazajian et al. 2009). *Right-hand side:* Redshift distribution for the LSST for $i < 24$, $i < 25$, $i < 27$ with $\sigma_i < 0.2$, and for galaxies with $S/N > 5$ in all bands, both for one and ten years of observations.

3.2.6. Apparent magnitude uncertainties for CFHTLS and the GOODS survey

An analytical expression similar to the one given by Eq. 2 does not exist for the CFHTLS and the GOODS data. The apparent magnitude uncertainties are estimated with algorithms and analysis techniques specific to these surveys and as such the relations in the previous section will not apply.

In the following, where simulations of photometric galaxy catalogs of both of these surveys will be carried out, the simulated uncertainties on the apparent magnitudes will be estimated directly from the survey data themselves. In this way, one can obtain the probability distribution of having σ_X given m_X . This will allow the assignment of σ_X , by randomly drawing the value according to this probability density function, given the value of m_X .

4. Method validation

4.1. GOODS

In order to validate the simulation scheme, we have performed a simulation of the GOODS WFI data⁵ and compared our results to the real photometric catalog used for the computation of the B -band luminosity functions reported in Dahlen et al. (2005).

The simulated photometric catalog corresponds to an effective solid angle of 1100 arcmin², that is equal to the area covered by the actual data catalog. The simulated redshift and absolute magnitude ranges are respectively $[0, 6]$ and $[-24, -13]$. The apparent magnitudes are computed for the WFI B -band and R -band. The apparent magnitude uncertainty is now given by the distribution of σ_X given m_X computed from the real data (see 3.2.6). If $m_{X,s}$ is the simulated apparent magnitude in any X -band, the uncertainty is randomly drawn from the distribution $\text{Prob}(\sigma_X | m_{X,s})$ found from the data. The observed apparent

⁵ Data from the Wide Field Imager (WFI) on the 2.2-m MPG/ESO telescope at La Silla.

magnitude and its uncertainty are then simulated as detailed in section 3.1.2.

Figure 4 shows the very good agreement of the galaxy number counts in the *R*-band between the simulation and the real data, except for the very faint galaxies at $R \geq 25$ for which the selection effect in the real data becomes important. The agreement is also very good for the *B*-band (not shown here). As displayed in Fig. 5, the distribution of colors from our GOODS simulation (black lines) are in reasonable agreement with the ones from the real data (red lines), for bright and faint galaxies. At bright magnitudes, the fitted luminosity function seems to predict a larger fraction of elliptical galaxies than the data. This feature probably comes from the SED templates and their linear interpolation. We have chosen to interpolate linearly between the SED templates with an equal number of steps between each template. With real galaxies it could easily be that this is not the case, e.g. perhaps there is not a uniform chance of a galaxy's SED lying between the E1 and the Sbc galaxy type, but instead there could be a higher chance it is more similar to the Sbc. This could explain our excess. However, the overall shape of the distributions indicates our simulated photometric catalog represents reality. This is expected because the luminosity functions were computed from the real data sample used for the comparison to the simulation.

4.2. CFHTLS

The different photo-*z* reconstruction methods were tested on real data, namely on galaxies observed both photometrically (from CFHTLS T0005) and spectroscopically (from either VVDS, DEEP2 or zCOSMOS surveys). We have followed the procedure described in detail by Coupon *et al.* (2009).

The CFHTLS T0005 public release contains a photometric catalog of objects observed in 5 bands (*u*, *g*, *r*, *i*, *z*) from the D1, D2, D3, W1, W3 and W4 fields. Among these, some galaxies are also spectroscopically observed by the VVDS, DEEP2 Redshift Survey and zCOSMOS surveys. The numbers of galaxies in the CFHTLS fields that were matched to spectroscopic observations are listed in Table 3. To perform the matching, the smallest angle between each galaxy in the CFHTLS catalog and a galaxy from the spectroscopic catalogs was computed. Only galaxies for which the angle is less than 0.7 arcsec (the order of the PSF) are grouped into the spectro-photometric catalog.

The spectroscopic redshift distribution of the spectro-photometric catalog is shown in Fig. 6.

A simulation of the CFHTLS data was also performed. This was to enable us to evaluate whether the statistical test described in section 5.2 can be calibrated with a simulation and then applied to real data, for which the photometric redshifts of galaxies are not known. Such a procedure would be useful when no spectroscopic sample is available to calibrate the prior probabilities or the likelihood ratio statistical test presented in section 5.2. The same argument stands for the neural network analysis described in section 7 which could also benefit from a simulated training sample.

Because the selection function of the spectro-photometric catalog is not based only on the detection threshold, the redshift and color distributions of the simulation and real data cannot be rigorously compared. The photometric selection criteria is not consistent over this data sample due to the differing selection of VVDS, DEEP2 and zCOSMOS, and therefore will not match the simulation. Since we also don't know how these selection criteria vary we cannot consistently compare the simulation to the data. However, the theoretical colors can be compared to the

Table 4. Values of the fit parameters of Eq. 3 and 4.

Spectral family:	<i>Early</i>	<i>Late</i>	<i>Starburst</i>
LSST			
α	2.97 ± 0.01	-1.57 ± 0.48	1.32 ± 0.00
z_0	0.00 ± 0.00	0.12 ± 0.02	0.00 ± 0.00
k_m	0.13 ± 0.00	0.08 ± 0.00	0.08 ± 0.00
f_t	0.22	0.41	...
k_t	0.27 ± 0.00	0.10 ± 0.00	...
CFHTLS			
α	3.51 ± 0.02	2.77 ± 0.01	1.89 ± 0.00
z_0	0.00 ± 0.00	0.18 ± 0.02	0.02 ± 0.01
k_m	0.13 ± 0.00	0.02 ± 0.01	0.08 ± 0.00
p_m	0	0.18 ± 0.02	0.02 ± 0.01
β	...	1.71 ± 0.08	3.12 ± 0.23
f_t	0.23	0.43	...
k_t	0.27 ± 0.00	0.11 ± 0.00	...

ones given by real data. To do so, the K-corrections are computed for each main galaxy type T_s , and this allows us to infer the colors as a function of the redshift. These are displayed in Fig. 7, for the *u* – *g* and *i* – *z* terms. The *g* – *r* and *r* – *i* exhibit a similar behavior. The data points and the theoretical curves are in agreement over the whole redshift range.

4.3. LSST forecasts

The simulated LSST sample considered here is the same as the one used in the companion paper Abate *et al. in prep.* It is generated with a solid angle of 7850 deg² and a redshift range of [0.1, 12]. This upper redshift limit is large enough to include all observable galaxies. The total number of galaxies is $\sim 8 \times 10^9$. For the purpose of this paper, namely the reconstruction of photometric redshifts, such a large number of galaxies is not necessary. Therefore the following analysis is done with a smaller subsample of galaxies, see section 6.2.

The expected cumulative number counts per unit of solid angle and per *i* band apparent magnitude for the LSST is shown on the left-hand side of Fig. 8, and is compared to the SDSS measurements made on the “Stripe 82” region (cf Abazajian *et al.* 2009). The LSST galaxy count is generally below that of SDSS, with a more pronounced effect at high and low end magnitudes ($m \sim 26$ and $m \sim 20$). There was no specific matching of the GOODS luminosity functions to the SDSS so we expect there to be some discrepancy due to comparing to a distribution that is different from the one it was drawn from.

The expected number of galaxies per unit of solid angle and per redshift is shown on the right-hand side of Fig. 8 for different cuts: *i* magnitude cuts of $i < 24$, $i < 25$, $i < 27$ with $\sigma_X < 0.2$ for all *X* bands, for both one (dotted lines) and ten years (solid lines) of observations. The LSST gold sample is defined as being all galaxies with $i < 25.3$ and is expected to produce high quality photometric redshifts.

The number of galaxies with S/N > 5 in all six bands at ten years is fairly small because these constraints are strong for the low acceptance *u* and *y* bands. Low S/N in the *u*-band is expected both from its shallower depth and from dropout galaxies




Fig. 9. Example of photometric computation for a simulated galaxy observed with the LSST in six bands at 5σ for ten years of observation. The 2D distributions correspond to the posterior probability density functions marginalized over the remaining parameter, and the 1D distributions correspond to the posterior probability density functions marginalized over the two remaining parameters. The top middle box corresponds to the value of the input parameters. On the top right-hand panel, the index *grid* denotes the parameters that maximize the 3D posterior probability density function on the grid, and on the middle right-hand panel the index *marg* denotes the parameters that maximize the 1D posterior probability density functions. The size of the grid cells has been reduced, and the z_p axis has been shortened compared to the size of the grid usually used to compute the likelihood function.




Fig. 10. Probability density function of the reduced variables $N_p(z_p)$, $N_p(T_p)$ and $g-r$. The black lines correspond to $P(\mu_i|G)$ and the red lines to $P(\mu_i|O)$.

at higher redshift where hydrogen absorption removes all flux blue-ward of the Lyman break, and non-detections above $z > 3$ are expected. The low S/N in the y-band is expected just from its shallower depth.

5. Enhanced template-fitting method

5.1. Maximum of the posterior probability density function

In this section, our template-fitting method for estimating photometric redshifts is presented. The algorithm follows the approach developed by Ilbert et al. (2006), Bolzonella et al. (2000) or Benitez (2000).

Basically, the template-fitting method consists in finding the photo- z z_p , the SED template T_p , the color excess $E(B-V)_p$ ⁶, and the SED normalization N that give the fluxes in each band that best fit to the observed values. Following Benitez (2000), the normalization parameter is marginalized over, so that the parameters of interest are given by the minimum of a χ^2 statistic whose expression will be given later in this section.

A Bayesian prior probability can be used to improve the photometric redshift reconstruction. It is defined as the probability of having a galaxy of redshift z and type T , given its apparent magnitude. It was introduced by Benitez (2000). Bayes' Theorem indicates that this probability can be expressed as the product of the probability of having a galaxy of type T given the apparent magnitude i : $P(T|i)$ times the probability of having a galaxy of redshift z , given the type and the apparent magnitude: $P(z|T, i)$. In other words

$$P(z, T|i) = P(z|T, i) \times P(T|i).$$

⁶ Subscript p refers to the “best-fit” parameters.




Fig. 11. Likelihood ratio distribution from the LSST simulation training sample. Logarithm of the probability density $P(L_R|G)$ in solid black and $P(L_R|O)$ in dashed red.

The two terms are well described by the functions:

$$P(T|i) = f_i e^{k_i(i-20)}, \quad (3)$$

and

$$P(z|T, i) \propto z^\alpha \exp\left[-\left(\frac{z}{z_m}\right)^\alpha\right], \quad (4)$$

$$z_m = z_0 + k_m(i - 16) + p_m(i - 16)^\beta.$$

Here, T represents the spectral family (broad type) instead of the spectral type (the exact SED), *i.e.* galaxies with spectral type lower than 5 belong to *Early*-type, those with spectral type between 6 and 25 belong to *Late*-type and the rest belong to *Starburst*-type. This parametrization follows a general model for galaxy number counts with redshift, and is improved to account for higher redshifts in CFHTLS. The parameters of $P(T|i)$ and $P(z|T, i)$ are found from fitting Eq. 3 and Eq. 4 to the simulated magnitude-redshift distributions for both the LSST and CFHTLS surveys. The value of the parameters in Eq. 3 and Eq. 4 are given by table 4. The fitted p_m parameter for the LSST is compatible with zero, while it is meaningful for CFHTLS (see table 4), it was therefore set to zero when representing the prior probability for the LSST simulations. There is no value for the parameters f_i and k_i for Starburst galaxies, whose probability distribution is set by the condition that the sum of probabilities over all galaxy types must be equal to 1.

When prior probabilities are taken into account, the χ^2 is extended and is defined as

$$\chi^2(z, T, E(B - V)) = \sum_X^{N_{band}} \left(\frac{F_{X,obs} - A F_{X,pr}(z, T, E(B - V))}{\sigma(F_{X,obs})} \right)^2 - 2 \ln(B) - 2 \ln(P(T|i) \times P(z|T, i)), \quad (5)$$

where $F_{X,obs}$ is the observed flux in the X -band; $F_{X,pr}(z, T, E(B - V))$ is the expected flux; N_{band} is equal to 5 for the CFHTLS, and is equal to 6 for the LSST; $\sigma(F_{X,obs})$ is the observed flux uncertainty. The terms A and B come from the analytical marginalization over the normalization of the SEDs; they are defined as follows:

$$A = \sum_{X=1}^{N_{band}} \frac{F_{X,obs} F_{X,pr}}{\sigma(F_{X,obs})} / \sum_{X=1}^{N_{band}} \frac{F_{X,pr} F_{X,pr}}{\sigma(F_{X,obs})}$$

$$B = \sum_{X=1}^{N_{band}} \frac{F_{X,pr} F_{X,pr}}{\sigma(F_{X,obs})}. \quad (6)$$

In the following, the 3D posterior pdf is defined as $\mathcal{L} = \exp[-\chi^2/2]$. It is computed for each galaxy on a 3D grid of $100 \times 25 \times 5$ nodes in the $(z, T, E(B - V))$ parameter space. The values of the parameters $z, T, E(B - V)$ lie in the intervals: $[0, 4.5], [0, 50], [0, 0.3]$ respectively. Since we are controlling the domain of possible parameter values to match the ranges used to make the simulation, we are probably reducing the number of possible degeneracies in $z, T, E(B - V)$ space. The SED library used for the photo- z reconstruction is the CWW+Kinney library described in Section 3.2.2. However, when considering the CFHTLS spectro-photometric data, the templates have been optimized following the technique developed by Ilbert et al. (2006); therefore naturally they match the data better.

The probability distribution is a function of 3 parameters; to derive the information on just one or two of the parameters we integrate the distribution over all values of the unwanted parameter(s) in a process called marginalization. The marginalized 2D probability density functions of the parameters $(z, T), (z, E(B - V)), (T, E(B - V))$ are computed in this manner, as well as the marginalized 1D probability density function of each parameter. Figure 9 shows an example of these probability density functions for a galaxy with true redshift $z_s = 0.16$, true type $T_s = 45$ (Starburst) and true excess color $E(B - V)_s = 0.24$. In many cases, the 3D posterior pdf is highly multimodal; minimizing the χ^2 with traditional algorithms such as MINUIT therefore often misses the global minimum. A scan of the parameter space is better suited to this application. Even a Markov Chain Monte Carlo (MCMC) method, which is usually more efficient than a simple scan, is not well suited for a multimodal 3D posterior pdf. Moreover, the production of the chains and their analysis in a 3D parameter space is slower than a scan. This example where $(z_p - z_s)/(1 + z_s) = 0.27$ corresponds to a catastrophic reconstruction. In this case, the parameters $(z_{pk}^{grid}, T_{pk}^{grid}, E(B - V)_{pk}^{grid})$ that maximize the 3D posterior pdf grid do not coincide with the ones that maximize the individual posterior probability density functions, namely the parameters $(z_{pk}^{marg}, T_{pk}^{marg}, E(B - V)_{pk}^{marg})$.

5.2. Statistical test and rejection of outliers

In this section, we outline a statistical test that aims at rejecting some of the outlier galaxies, for which $|z_p - z_s|/(1 + z_s) > 0.15$. It is based on the characteristics of the 1D posterior probability density functions $P(z_{pk})$, $P(T_{pk})$ and $P(E(B - V)_{pk})$. The test is calibrated with a training sample, for which the true redshift is known (or the spectroscopic redshift in the case of real data). In the following, the LSST simulation for ten years of observation is considered in order to illustrate the method.

5.2.1. The probability density function characteristics

The variables considered in order to establish the statistical test are the following:

- The number of peaks in the marginalized 1D posterior probability density functions denoted by $N_{pk}(\theta)$, where θ is either z, T or $E(B - V)$.
- When $N_{pk} > 1$, the logarithm of the ratio between the height of the secondary peak over the primary peak in the 1D posterior probability density functions, denoted $R_L(\theta)$.




Fig. 12. From the LSST simulation training sample. Top panel: evolution of the rejection vs. L_R . Bottom left-hand panel: evolution of the acceptance vs. the rejection. Bottom right-hand panel: evolution of the acceptance vs. L_R .

- When $N_{pk} > 1$, the ratio of the probability associated with the secondary peak over the probability associated to the primary peak in the 1D posterior probability density functions, denoted by $R_{pk}(\theta)$. The probability is defined as the integral between two minima either side.
- The absolute difference between the value of z_{pk} and z_{pk}^{marg} , denoted by $D_{pk} = |z_{pk} - z_{pk}^{marg}|$, where z_{pk}^{marg} is the redshift that maximizes the posterior probability density function $P(z)$.
- The maximum value of $\log(\mathcal{L})$.
- The colors $\mathcal{C} = (u-g, g-r, r-i, i-z)$ in the case of CFHTLS, and with an extra $z-y$ term in the case of the LSST.

We denote by O the galaxies that are considered as outliers and G the galaxies for which the redshift is well reconstructed in the following way:

- O : $|z_p - z_s|/(1 + z_s) > 0.15$
- G : $|z_p - z_s|/(1 + z_s) < 0.15$

We define the set of variables defined in the list above are denoted by the vector μ . From a given training sample, we compute the distributions $P(\mu_i|O)$ and $P(\mu_i|G)$. For convenience we adopt reduced variables that are renormalized to lie between 0 and 1. Distributions of some of the reduced variables are plotted in Figs. 10. It is clear that the distributions $P(\mu_i|G)$ and $P(\mu_i|O)$ are different. The probability that an outlier galaxy O presents more than three peaks in its posterior probability density function $P(z_p)$ is larger than for a well reconstructed galaxy G . A combination of these different pieces of information leads to an efficient test to discriminate between good and catastrophic reconstructions.

5.2.2. Likelihood ratio definition

In order to combine the information contained in the densities $P(\mu_i|G)$ and $P(\mu_i|O)$, we define the likelihood ratio variable L_R :

$$L_R(\mu) = \frac{P(\mu|G)}{P(\mu|G) + P(\mu|O)},$$




Fig. 13. From the LSST simulation training sample. 2D distribution of $(z_p - z_s)/(1 + z_s)$ as a function of the threshold $L_{R,c}$. In each bin in $L_{R,c}$, the distribution of $(z_p - z_s)/(1 + z_s)$ is normalized. We find $L_{R,c} > 0.98$ to be a good choice.




Fig. 14. Histogram of the number of galaxies from the CFHTLS sample with $L_R \geq L_{R,c}$ as a function of $L_{R,c}$. The black curve has been obtained with densities $P(\mu|G)$ and $P(\mu|O)$ computed from the CFHTLS data themselves whereas the red curve relies on densities obtained from the CFHTLS simulation.

where

$$P(\mu|G) = \prod_{i=1}^{N_\mu} P(\mu_i|G),$$

$$P(\mu|O) = \prod_{i=1}^{N_\mu} P(\mu_i|O),$$

where N_μ is the number of components of μ . Here, the variables μ_i are assumed to be independent. The correlation matrix of the μ_i 's shows indeed a low correlation between the parameters. We approximate the two probabilities $P(G|\mu)$ and $P(O|\mu)$ as the product of $P(\mu|G)$ and $P(\mu|O)$, neglecting the correlations, as our aim is just to define a variable for discriminating the two possibilities. The probability density functions $P(L_R|G)$ and $P(L_R|O)$ are computed from the training sample, and are displayed in Fig. 11. The results shown here are from the LSST simulation for ten years of observations with $m_X < m_{5,X}$. As expected, the two distributions are very different.

The quality of a discrimination test, such as $L_R > L_{R,c}$, can be quantified by the acceptance Acc and rejection Rej rates:

$$Acc(L_R) = \int_{L_R}^1 P(L'_R|G)dL'_R,$$

$$Rej(L_R) = \int_0^{L_R} P(L'_R|O)dL'_R.$$

The evolution of Acc and Rej as functions of L_R and of Acc as a function of Rej is displayed in Fig. 12. The larger the difference between the curve Acc vs. Rej and the curve $Acc = 1 - Rej$, the higher the rejection power. Figure 12 shows that the method should work because the solid line lies far from the diagonal dotted line in the bottom left panel. A high value of L_R is necessary to discard outliers, however it should be chosen so a minimum of well-reconstructed galaxies are removed. The distribution of $(z_p - z_s)/(1 + z_s)$ as a function of the cut on L_R , namely $L_{R,c}$, is displayed in Fig. 13, which is a visual illustration of the L_R cut effectiveness. The plots in Figs. 15 & 16, discussed below, show that there is a significant improvement when a cut on L_R is applied.

6. Photo-z performance with template-fitting

In the following sections, the ability of the statistical test, based on the L_R variable, to construct a *robust sample* of galaxies with well reconstructed redshifts is investigated in more detail, both for the CFHTLS spectro-photometric data and the LSST simulation. The efficiency of the photo-z reconstruction is quantified by studying the distribution of $(z_p - z_s)/(1 + z_s)$, through the:

- **bias**: median that splits the sorted distribution in two equal samples.
- **rms**: the interquartile range (IQR)⁷. If the distribution is Gaussian, it is approximately equal to 1.35σ where σ is the standard deviation.
- **η** : the percentage of outliers for which $|z_p - z_s|/(1 + z_s) > 0.15$.

Table 2 gives the LSST requirements for these values.

6.1. Results for CFHTLS

In this section, the reconstruction of the photometric redshifts, and the consequences of the selection on L_R for the spectro-photometric data of CFHTLS, are presented in order to validate the method. Two cases are considered:

- **Case A**: The distributions $P(\mu|G)$ and $P(\mu|O)$ are computed from the data themselves.
- **Case B**: The distributions $P(\mu|G)$ and $P(\mu|O)$ are computed from a simulation of the CFHTLS data, as explained in section 4.2.

In both cases, the photo-z is computed from the real CFHTLS data. In order to compare the effect of a selection on L_R , in particular to keep the same total number of galaxies for a particular selection, different values of $L_{R,c}$ have to be chosen for Case A and Case B. For a given number of galaxies left in the sample, the values of $L_{R,c}$ for both cases can be obtained from Fig. 14.

Figure 15 shows the evolution of the number of galaxies, the bias, rms, and the parameter η as functions of z_s , for three values

of $L_{R,c}$ for Case A (solid lines) and B (dashed lines). The photometric redshifts are clearly improved if a large enough value of L_R is applied (i.e. $L_R > 0.9$), even in Case B. However, within the galaxies that are removed, there are many well reconstructed galaxies.

In the top panel of Fig 15 larger L_R values remove more galaxies from the lowest redshift bins, indicating that probably $L_R > 0.6$ is required to remove outlier galaxies that have been assigned a low redshift due to e.g. confusion between the Lyman and Balmer breaks. In the second panel the L_R selection does not have a very strong effect on the bias, probably because our definition of bias is not heavily affected by outliers. The rms on the other hand, is improved by up to a factor of 3 at higher redshifts, and similarly the catastrophic outlier rate is also improved by up to a factor of 3 at higher redshifts. However, the efficiency of the L_R selection on the CFHTLS data is significantly below that on the LSST simulated data (see the following subsection). The CFHTLS spectro-photometric sample used here will predominantly contain bright, well measured galaxies, because these are the kind generally selected for spectroscopic follow-up. Therefore we expect there to be far fewer galaxies in this sample that have outlier photometric redshifts, and as such the L_R calibration is less robust, leading to the removal of too many well reconstructed galaxies.

In Fig. 6 we see that the spectroscopic sample redshift coverage barely extends beyond $z > 1.4$. This means that when the L_R selection was calibrated on the CFHTLS data, the sample was missing outlier galaxies with high redshifts that could be estimated erroneously to be at low redshifts, e.g. those subject to the degeneracy causing the Lyman break to be confused for the 4000 Å break. The LSST simulated data will contain these degeneracies, however it would be good to test the L_R selection method on real data out to higher redshifts in the future.

The agreement between the solid (Case A) and dashed (Case B) lines in Fig. 15 show that, if no spectroscopic survey is available, a simulation that reproduces the colors of the photometric catalog can be used to calibrate the likelihood ratio test. The model used to compute the photo-z, that is to say the CWW+Kinney library, the Calzetti and Cardelli extinction laws, correctly accounts for the data.

6.2. Results for LSST

We use a total of 50 million galaxies in our simulated catalog. This catalog is divided into 5 different sets. Each set is separated into a test sample (2 million galaxies) and an analysis sample (8 million galaxies). In each set, the statistical test is performed on "observed" galaxies within the test sample, then the densities $P(\mu|G)$ and $P(\mu|O)$ are used to compute the value of L_R for "observed" galaxies in the analysis sample. Performing the reconstruction on the 5 independent sets give us a measure of the fluctuation from a set to another and thus an estimate of the error on our reconstruction parameters. We performed the same analysis with 10 sets of twice less galaxies and measured the fluctuations to be very similar.

6.2.1. Observation in six bands

To test the method with best photometric quality we require each galaxy to be "observed" in each band with good precision $m_X < m_{5,X}$. This requirement leaves us with about 125 000 galaxies in the test sample and 500 000 in the analysis sample.

⁷ The interquartile range is the interval spanning the second and third quartiles.




Fig. 15. CFHTLS spectro-photometric data. *Top*: evolution of the fraction of galaxies and the bias with z_s . *Bottom*: evolution of the rms and η as a function of z_s . With Case A (calibrated with data) shown by the solid lines and Case B (calibrated with simulation) by the dashed lines. The different values of $L_{R,c}$ for Case A are reported in the legend, and for Case B in parentheses in the legend.




Fig. 16. Distribution of $z_p - z_s$ versus z_s for a simulated LSST catalog, for all galaxies (left) and for galaxies with likelihood ratio L_R greater than 0.98 (right).

Figure 16 shows the 2D distributions from the LSST simulation of $z_p - z_s$ as a function of z_s for all the galaxies in the sample compared to the same distribution after performing an L_R selection. It is clear that selecting on the likelihood ratio enhances the photometric redshift purity of the sample.

Figure 17 (the same as Fig. 15 for CFHTLS) shows the evolution with z_s of the number of galaxies retained in the LSST sample, for each of the parameters listed above (bias, rms, η).

This indicates the quality of the photo-z, for different values of $L_{R,c}$.

The LSST specifications on the bias and rms (see table 2) are fulfilled up to $z_s = 1.5$ with only a low value of $L_{R,c} > 0.6$. For redshifts greater than 1.5, a higher value of $L_{R,c}$ is required in order to reach the expected accuracy. There are two main reasons for this. Firstly, only a small percentage of the galaxies with $z_s > 1.5$ are used to calibrate the densities $P(\mu|G)$ and $P(\mu|O)$,




Fig. 17. LSST L_R selection. *Top*: evolution of the fraction of galaxies and the bias with z_s . *Bottom*: evolution of the rms and η as a function of z_s . The thick black lines represent the LSST requirements given in table 2. Ten years of observations with the LSST is assumed. The values of $L_{R,c}$ are reported in the legend.

therefore the high-redshift galaxies do not have much weight in the calibration test. Secondly, the ratio between the height of the distribution $P(L_R|G)$ at $L_R = 0$ and at $L_R = 1$ tends to increase with the redshift, meaning that the purity of the test is degraded. Finally, only a very low value of $L_{R,c} > 0.2$ is needed to enable η to meet LSST specifications for $z_s > 2.2$.

The effect of a selection on the likelihood ratio L_R can be compared to the effect of a selection on the apparent magnitude in the i -band, as shown in Fig. 18. The increase in the i -magnitude selection efficiency at large z shown in Fig. 18 (top) is due to the value of i_{cut} approaching the detection threshold. Performing a selection on a quantity other than magnitude, such as the likelihood ratio, ensures that “well measured” but faint galaxies are still included in the sample.

Of course requiring observation in six bands will exclude the u -band drop out galaxies at high redshift, so the photo- z performance at these redshifts will be greatly affected by this requirement. The next section investigates the photo- z performance when observations are required in less than six bands.

6.2.2. Observation in five bands (and less)

The previous subsection demonstrated our results for good photometric data with $m_X < m_{5,X}$ in all six bands of the LSST. In order to detect more galaxies and extend our reconstruction at higher redshift, we release the constraint on the number N_{m5} of “well observed” bands having $m_X < m_{5,X}$. Both test and analy-

Table 5. LSST number of galaxies, comparison between the selection on L_R and the selection on i band magnitude. Observations were required in at least 5 bands.

	Total	$0 \leq z < 1$	$1 \leq z < 2$	$2 \leq z < 3$
$L_{Rc} = 0$	893123	537016	326571	29536
$L_{Rc} = 0.98$	426667	311997	110960	3710
$i < 24$	314461	263207	50084	1171

sis sample are made of galaxies with $m_X < m_{5,X}$ in at least N_{m5} bands. We decreased N_{m5} from 6 to 5, 4 and 3 and performed similar analysis to what was presented in the previous subsection. The comparison of the results indicates that the selection with $N_{m5} = 5$ gives the best results. As we can see in Fig. 19, lowering N_{m5} from 6 to 5 greatly increases the number of galaxies we keep in our sample without significantly degrading the reconstruction performance. The gain in the number of galaxies is presented in table 6 and increases with redshift as expected.

For galaxies “observed” in only 5 bands, the band X which has $m_X > m_{5,X}$ or is not observed at all (noise level) is the u band in 95% of the cases.

When requiring less than 5 bands, the results are worse or similar: in order to reconstruct decent photometric redshifts in this case we need to apply such a large value of L_R for the selec-




Fig. 18. LSST i magnitude selection. *Top*: evolution of the fraction of galaxies and the bias with z_s . *Bottom*: evolution of the rms and η as a function of z_s . The thick black lines represent the LSST requirements given in table 2. Ten years of observations with the LSST is assumed. The values of the i -magnitude cuts are reported in the legend.

Table 6. LSST number of galaxies ($LR_c = 0.98$)

	Total	$0 \leq z < 1$	$1 \leq z < 2$	$2 \leq z < 3$
$N_{m5} = 6$	296715	235059	60818	838
$N_{m5} = 5$	426667	311997	110960	3710
$N_{m5} = 4$	533363	375966	149748	7649
$N_{m5} = 3$	534307	376772	149715	7819

tion that we reject nearly all the galaxies gained from the weaker requirement, and even discard well-measured galaxies.

Figure 20 shows the comparison of number of galaxies, and photo- z performance (rms, bias, η) for a L_R selected sample ($L_R > 0.98$) and a magnitude selected sample ($i < 24$). While both samples satisfy the LSST science requirement given in table 2, the L_R selection is more efficient, since it retains a significantly larger number of galaxies, specifically for $z > 1$ (see table 5). We do not present a comparison with a sample selected by a magnitude cut of $i < 25.3$ since it would not satisfy the LSST photo- z requirements according to our simulation and photo- z reconstruction (as can be seen in Fig. 18).

7. Photo- z performance with neural network

It has been shown using the public code ANNz by Collister & Lahav (2004) that the photometric redshifts can be correctly es-

timated via a neural network. This technique, among other empirical methods, requires a spectroscopic sample for which the apparent magnitudes and the spectroscopic redshifts are known.

The Toolkit for Multivariate Analysis (Hoecker et al. 2007, TMVA) provides a ROOT-integrated environment for the processing, parallel evaluation and application of multivariate classification and multivariate regression techniques. All techniques in TMVA belong to the family of supervised learning algorithms. They make use of training events, for which the desired output is known, to determine the mapping function that either describes a decision boundary or an approximation of the underlying functional behavior defining the target value. The mapping function can contain various degrees of approximations and may be a single global function, or a set of local models. Among artificial neural networks, many other algorithms, such as boosted decision trees or support vector machine are available. An advantage of TMVA lies in the fact that different algorithms can be tested at the same time in a very user-friendly way.

7.1. Method

The MultiLayer Perceptron (MLP) neural network principle is quite simple. It builds up a linear function that maps the observables to the target variables, in our case, the redshifts. The coefficients of the function, namely the weights, are such that they minimize the error function that is the sum over all galaxies in the sample of the difference between the output of the network




Fig. 19. LSST reconstruction for different N_{m5} requirement. *Top:* evolution of the fraction of galaxies and the bias with z_s . *Bottom:* evolution of the rms and η as a function of z_s . The thick black lines represent the LSST requirements given in table 2. Ten years of observations with the LSST is assumed. The values of N_{m5} and L_{Rc} are reported in the legend.

and the true value of the target. Two samples of galaxies are necessary, the training and the test sample. The latter is used to test the convergence of the network and to evaluate its performance. It usually prevents over-training of the network, which may arise when the network learns the particular feature of the training sample.

A neural network is built with layers and nodes. There are at least two layers, one for the input observables $\mathbf{x}$ and one for the target z_{MLP} . Each node of a layer is related to all nodes from the previous layer, with a weight w , which is the coefficient associated with the activation function A of each connection. The value y_j^{i+1} of the node k from the layer $i + 1$ is related to the values of all y_j^i from the layer i :

$$y_k^{i+1} = A \left(\sum_{j=1}^n w_{jk}^i y_j^i \right),$$

where n is the number of neurons in the i layer. For the purpose of this paper, the activation function A is a sigmoid. As an example, in the case where there is only one intermediate layer, the photometric redshift of the g^{th} galaxy is estimated as follows:

$$z_{MLP,g} = \sum_{j=1}^n A \left(\sum_{i=1}^{n_{band}} w_{ij}^1 x_{g,i} \right) \times w_{j1}^2.$$

The error function is simply defined by

$$E(\mathbf{w}) = \frac{1}{2} \sum_{g=1}^{n_{train}} E_g(\mathbf{x}_g, \mathbf{w}),$$

with

$$E_g(\mathbf{x}_g, \mathbf{w}) = z_{MLP,g}(\mathbf{x}_g, \mathbf{w}) - z_{s,g},$$

where n_{train} is the number of galaxies in the training sample and g denotes the g^{th} galaxy. At the first iteration, the weights have random values. The gradient descent method, which consists of modifying the weight value according to the derivative of E with respect to the weight, is used to minimize E . For example, after one iteration, we have:

$$w_{j1}^2 \rightarrow w_{j1}^2 + \Delta w_{j1}^2,$$

$$\Delta w_{j1}^2 = -\alpha \sum_{g=1}^{n_{train}} \frac{\partial E_g}{\partial w_{j1}^2}.$$

The parameter α is the learning rate, and has to be determined for each specific case. It must not be too large, otherwise the steps are so large that the minimum of E is never reached. It must not be too small either, otherwise too many iterations are required. The testing sample is used as a convergence and performance test. Indeed, the errors decrease with the number of iterations




Fig. 20. LSST comparison between $L_{Rc} = 0.98$ and $i < 24$ cuts. *Top*: evolution of the fraction of galaxies and the bias with z_s . *Bottom*: evolution of the rms and η as a function of z_s . The thick black lines represent the LSST requirements given in table 2. Ten years of observations with the LSST is assumed.




Fig. 21. Comparison between the template-fitting method and the neural network for CFHTLS data. The bias, the rms of the distribution of $\Delta z/(1+z_s)$, and the parameter η are displayed as functions of the true redshift for the CFHTLS data. Data points are reported only if the number of galaxies in the sample is greater than ten.

in the training sample, but reach a constant value on the testing sample. Weights are finally kept when the errors on the testing sample reach a constant value.

For this non-exhaustive study, we have chosen the observables $\mathbf{x} = (\mathbf{m}, \sigma(\mathbf{m}))$, and two layers of ten nodes each. In Fig. 21, the bias, rms and the outlier rate η are compared for the template-fitting method and for the neural network. It is clear that the outlier rate is much smaller at all redshifts when the photo- z is estimated from the neural network. The dispersion of the photometric redshifts is also smaller for the neural network when compared to the template-fitting method. These

characteristics are expected because the training sample is very similar to the test sample, whereas the template-fitting method uses only a small amount of prior information. Moreover, in the template-fitting method, the apparent magnitudes are fitted with a model, with a SED templates, whereas to run the neural network no theoretical model have to be assumed. However, these attributes may be reversed if the two samples are different, as will be illustrated in Section 7.3. The over-estimation of photo- z at low redshifts, and under-estimation at high redshifts, shown by the downward slope in the bias, can be attributed to *attenuation bias*. This is the effect of the measurement errors in the

observed fluxes resulting in the measured slope of the linear regression to be under-estimated on average; see Freeman *et al.* (2009) for a full discussion of this bias. We note that the photo- z bias obtained from the template-fitting method has an opposite sign, and is of the same amplitude, to that obtained from the neural network method. Since we have reason to expect the neural network has a downward slope in the bias, this indicates that the two estimators can be used complementarily. This is investigated in the next subsection.

7.2. Results for CFHTLS

Here, a possible combination of photo- z estimators (from the template-fitting method on the one hand and from the neural network on the other hand) is outlined. Even if the fraction of outlier galaxies is smaller with a neural network method than for the template-fitting method, using only the neural network to estimate the photo- z appears not to be sufficient to reach the stringent photo- z requirements for the LSST, especially when the spectroscopic sample is limited. In fact, neural networks seem to produce a photo- z reconstruction slightly biased at both ends of the range (see Fig. 21). This is due to the galaxy under-sampling at low and very high redshifts in the training sample. In case spectroscopic redshifts are available, it is therefore worth combining both estimators.

With the CFHTLS data, there is a correlation between $z_p - z_{MLP}$ and $z_p - z_s$, where z_p is the photo- z estimated with the template-fitting method, as displayed in Fig. 22. This correlation could be used to remove some of the outlier galaxies for which the difference between z_p and z_s is large, for example, by removing galaxies with around $|z_p - z_{MLP}| \geq 0.3$. This correlation appears because, in the case where the neural network is well trained, as is the case here, the photo- z is well estimated and z_{MLP} becomes a good proxy for z_s .

One can see an example of the impact of using both estimators z_p and z_{MLP} in Fig. 23. The distribution of $z_p - z_s$ is plotted for three cases: with $L_R > 0.9$ only, with $|z_p - z_{MLP}| < 0.3$ only and with both cuts. There are fewer outlier galaxies from the first to the third case. This shows that if the training sample is representative of the photometric catalog, neural networks have the capability to tag galaxies with an outlier template-fitted photo- z . However this will be difficult to achieve in practice because the training sample is biased in favor of bright, low redshift galaxies, that are in majority selected for spectroscopic observations. Selecting with both variables improves the photo- z estimation compared to a cut only on $|z_p - z_{MLP}|$ but to a lesser extent.

7.3. Results for the LSST

For the LSST simulation the network was composed of 2 layers of 12 nodes each, the training sample was composed of 10 000 galaxies and the testing sample of 20 000 galaxies. We found that increasing the size of the training sample above 10 000 showed no improvement in the precision of the training. We attribute this to the regularity of the simulation: the galaxies were drawn from a finite number of template SEDs, therefore as soon as the sample represents all the galaxy types in the simulation, adding more galaxies does not help in populating the parameter space any longer.

A scatter plot of photo- z versus spectroscopic redshift is shown on the top panel of Fig. 24. The black points show the results from the template-fitting method, where a selection of $L_R > 0.98$ was applied, and the red points show the results




Fig. 22. 2D histogram of $z_p - z_{MLP}$ vs. $z_p - z_s$ for the test sample of the CFHTLS data.




Fig. 23. Normalized histograms of $z_p - z_s$ with $L_R > 0.9$ (black curve), $|z_{MLP} - z_p| < 0.3$ (red curve) and both cuts (blue curve) for the CFHTLS data.

from the neural network described above. The plot compares the photo- z performance of the neural network method and the template-fitting method on the simulated LSST data. Similar to Singal *et al.* (2011), we find that the neural network results in fewer outliers, although it has a larger rms for “well measured” galaxies than the template-fitting method.

In the bottom panel of Fig. 24 the correlation between $z_p - z_{MLP}$ and $z_p - z_s$ is shown. Here the correlation between both estimators is less useful for identifying outliers than it was for CFHTLS. This is presumably due to both the simulation and the fit being performed with the same set of galaxy template SEDs. This should significantly reduce the fraction of outliers compared to a case where the templates used to estimate z_p do not correctly represent the real galaxies. For example, removing some of the templates from the z_p fit reduces the photo- z quality, as demonstrated in Benitez (2000). Therefore the existence of a strong correlation between $z_p - z_{MLP}$ and $z_p - z_s$ may be useful in diagnosing and mitigating problems with the SED template set.

It is difficult to obtain a spectroscopic sample of galaxies that is truly representative of the photometric sample, in terms of redshifts and galaxy types (Cunha *et al.* 2012). For example, in the case of the LSST, the survey will be so deep that spectroscopic redshifts will be very hard to measure for the majority




Fig. 24. Top panel: z_p vs. z_s with the template-fitting method in black (selection with $L_R > 0.98$), and with the neural network in red, for an LSST simulation of ten years of observations. Bottom panel: 2D histogram of $z_{MLP} - z_p$ as a function of $z_p - z_s$.

of faint galaxies or those within the “redshift desert”. Here, we briefly investigate the effect of having the spectroscopic redshift distribution of the training sample biased with respect to the full photometric sample.

The fact that the distribution of redshifts in the spectroscopic sample is different from the underlying distribution is often (confusingly) termed “redshift bias”. The consequence of this bias can be seen by modifying the efficiency of detection as a function of the redshift. The efficiency function is chosen to be

$$\epsilon(z) = 1 - 1 / \left(1 + e^{-\left(\frac{z-1.2}{0.1} \right)} \right) \quad (7)$$

and it is plotted in Fig. 25 (inset). This efficiency function is then used to bias the training sample and the test sample, in order to compute new network weight coefficients. The photometric redshifts for another unbiased sample are then computed using these weights.

The scatter plot of $z_{MLP} - z_s$ as a function of z_s is shown in Fig. 25. We find that the photometric redshifts are quite well estimated as long as $\epsilon \geq 0.2$. This figure shows qualitatively that a bias in the training sample has a major impact on the photo- z reconstruction performance by the neural network, at least with the training method used here.




Fig. 25. $z_{MLP} - z_s$ as a function z_s for the ten years of observations of the LSST. The curve in the inset shows the efficiency function ϵ as a function of the redshift, as it is used on the training sample to force a bias in the redshift selection.

8. Discussion and future work

In regard to simulations undertaken here, there are a number of simplifications that will be reconsidered in future work. We discuss briefly some of these here.

- *Point source photometric errors:* We have assumed photometric errors based on estimates valid for point sources, and since galaxies are extended sources we expect the errors to be larger in practice. We made an independent estimate of the photometric errors expected for the LSST, including the error degradation due to extended sources. We found that for the median expected seeing, the photometric error scales as $\sigma_F/F = \theta/0.7$ where σ_F is the error on the flux F , and θ is the size of the galaxy in arcseconds. The next round of simulations will therefore include a prescription for simulating galaxy sizes in order to improve our simulation of photometric errors. We will also compare our simple prescription to results obtained from the LSST image simulator (ImSim).
- *Galactic extinction (Milky Way):* Our current simulations effectively assume that i) Galactic extinction has been exactly corrected for, and ii) our samples of galaxies are all drawn from a direction of extremely low and uniform Galactic extinction. In practice, there will be a contribution to the photometric errors due to the imperfect correction of the Galactic extinction, and this error will vary in a correlated way across the sky. More problematically, the extinction has the effect of decreasing the depth of the survey as a function of position on the sky. To account for these effects we will construct a mapping between the coordinate system of our simulation, and Galactic coordinates in order to apply the Galactic extinction in the direction to every galaxy in our simulation. We can use the errors in those Galactic extinction values to propagate an error to the simulated photometry.
- *Star contamination:* M-stars have extremely similar colors to early type galaxies and as such can easily slip into photometric galaxy samples. Taking an estimate for the expected LSST star-galaxy separation quality, we plan to contaminate our catalog with stars. This could have an important effect by biasing the clustering signal of galaxies since the contamination will be worse closer to the Galactic plane.

- *Enhanced SEDs*: Our current simulations are probably more prone to problems with degeneracies in color space because we use a uniform interpolation between the main type SEDs. This may lead to poorer photometric redshifts than would be expected in reality, since galaxies might not exhibit such a continuous variation in SED type. In the future we plan to implement a more realistic interpolation scheme as well as the use of more complete template libraries, such as synthetic spectral libraries.

Further improvements to the photo- z determination can be made by the use of angular cross-correlation between objects in the photo- z sample, and objects with spectroscopic redshifts, located in the same area of the sky (see Matthews & Newman (2010) and Matthews & Newman (2011)). This cross-correlation will help to characterize the redshift distribution of the photometric sample, even though the spectroscopic sample may be incomplete or otherwise not closely resemble the photometric sample.

9. Conclusion

We have developed a set of software tools to generate mock galaxy catalogs (as observed by the LSST or other photometric surveys) and to compute photometric redshifts and study the corresponding redshift reconstruction performance.

The validity of these mock galaxy catalogs was carefully investigated (see section 4). We have shown that our simulation reproduces the photometric properties of the GOODS and CFHTLS observations well, in particular the number count, magnitude and color distributions. We developed an enhanced template-fitting method for estimating the photometric redshifts which involves a likelihood ratio statistical test using the posterior probability functions of the fitted photo- z parameters (z , galaxy type, extinction ...) and the galaxy colors.

This method was applied both to the CFHTLS data and to the LSST simulation to derive photo- z performance, which was compared to the photo- z reconstruction using a multilayer perceptron (MLP) neural network. We have shown how results from our template fitting method and from the neural network might be combined to provide a galaxy sample of enriched objects with reliable photo- z measurements.

We find our enhanced template method produces photometric redshifts that are both realistic, and meet LSST science requirements when the galaxy sample is selected using the likelihood ratio statistical test. We have shown that a selection based on the likelihood ratio test performs better than a simple selection based on apparent magnitude, as it retains a significantly larger number of galaxies, especially at large redshifts ($z \gtrsim 1$), for a comparable photo- z quality.

We confirm that LSST requirements for photo- z determination, i.e. (2 – 5)% dispersion on the photo- z estimate, with less than $\sim 10\%$ outliers can be met, up to redshift $z \lesssim 2.5$. A number of enhancements for the mock galaxy catalog generation and photo- z reconstruction have been identified and were discussed in section 8.

The photo- z computation presented here is designed for a full BAO simulation aiming at forecasting the precision on the reconstruction of the dark energy equation-of-state parameter. This will be presented in a companion paper (Abate *et al.*, *in prep.*).

Acknowledgements. Thanks to Pierre Astier and Delphine Hardin for their advice on CFHTLS data, to Tomas Dahlen for his help on the simulation of the

GOODS mock galaxy catalog and Eric Gawiser for his help with the IGM calculation. We also thank the LSST internal reviewers: Patricia Burchat, Andy Connolly and Sam Schmidt for their constructive criticism of the original paper draft.

References

- Abazajian, K. N. *et al.* 2009, *Astrophys.J.Suppl.*, 182, 543
 Baum, W. A. 1962, in *IAU Symposium*, ed. M. G. C., Vol. 15, 390
 Benítez, N. 2000, *Astrophys. J.*, 536, 571
 Bolzonella, M., Miralles, J.-M., & Pello', R. 2000, *Astron. Astrophys.*, 363, 476
 Bruzual, G. & Charlot, S. 2003, *MNRAS*, 344, 1000
 Budavári, T., Szalay, A. S., Connolly, A. J., Csabai, I., & Dickinson, M. 2000, *AJ*, 120, 1588
 Calzetti, D. *et al.* 2000, *Astrophys. J.*, 533, 682
 Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *Astrophys. J.*, 345, 245
 Clayton, G. C., Gordon, K. D., Salama, F., *et al.* 2003, *ApJ*, 592, 947
 Coleman, G. D., Wu, C., & Weedman, D. W. 1980, *ApJS*, 43, 393
 Collister, A. A. & Lahav, O. 2004, *Publ. Astron. Soc. Pac.*, 116, 345
 Connolly, A., Csabai, I., Szalay, A., *et al.* 1995, *Astron.J.*, 110, 2655
 Coupon, J., Ilbert, O., Kilbinger, M., *et al.* 2009, *A&A*, 500, 981
 Cunha, C. E., Huterer, D., Busha, M. T., & Wechsler, R. H. 2012, *MNRAS*, 423, 909
 Dahlen, T. *et al.* 2005, *Astrophys. J.*, 631, 126
 Freeman, P. E., Newman, J. A., Lee, A. B., Richards, J. W., & Schafer, C. M. 2009, *MNRAS*, 398, 2012
 Garilli, B., Le Fèvre, O., Guzzo, L., *et al.* 2008, *A&A*, 486, 683
 Hoecker, A., Speckmayer, P., Stelzer, J., *et al.* 2007, *ArXiv Physics e-prints*
 Hogg, D. W. 1999, *ArXiv Astrophysics e-prints*
 Ilbert, O., Arnouts, S., McCracken, H. J., *et al.* 2006, *A&A*, 457, 841
 Ivezić, Z., Tyson, J. A., Acosta, E., *et al.* 2008, *ArXiv e-prints*
 Kinney, A. L., Calzetti, D., Bohlin, R. C., *et al.* 1996, *ApJ*, 467, 38
 Le Fevre, O., Guzzo, L., Meneux, B., *et al.* 2004, *Astron.Astrophys.*
 Lilly, S. *et al.* 2007, *Astrophys.J.Suppl.*, 172, 70
 LSST Science Collaboration. 2009, *ArXiv e-prints*
 Madau, P. 1995, *ApJ*, 441, 18
 Matthews, D. J. & Newman, J. A. 2010, *ApJ*, 721, 456
 Matthews, D. J. & Newman, J. A. 2011, *ArXiv e-prints*
 Newman, J. A., Cooper, M. C., Davis, M., *et al.* 2012, *ArXiv e-prints*
 Petry, C. E., Miller, M., Cook, K. H., *et al.* 2012, in *American Astronomical Society Meeting Abstracts*, Vol. 219, American Astronomical Society Meeting Abstracts #219, 156.09
 Puschell, J., Owen, F., & Laing, R. 1982, in *IAU Symposium*, Vol. 97, *Extragalactic Radio Sources*, ed. D. S. Heeschen & C. M. Wade, 423
 Sheldon, E. S., Cunha, C. E., Mandelbaum, R., Brinkmann, J., & Weaver, B. A. 2012, *ApJS*, 201, 32
 Singal, J., Shmakova, M., Gerke, B., Griffith, R. L., & Lotz, J. 2011, *PASP*, 123, 615
 Vanzella, E., Cristiani, S., Fontana, A., *et al.* 2004, *Astron.Astrophys.*, 423, 761